





Survey of back and neck pain in South African ophthalmologists



Authors:

Craig D. Anderson¹ 
Aubrey Makgotloe¹ 
Karen Koetsie² 
André Rose³ 

Affiliations:

¹Division of Ophthalmology, Department of Neurosciences, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

²Department of Ophthalmology, Yeovil Hospital NHS Foundation Trust, Yeovil, United Kingdom

³Centre for Health Systems Research & Development, University of the Free State, Bloemfontein, South Africa

Corresponding author:

Craig Anderson,
craigbyte@gmail.com

Dates:

Received: 24 Sept. 2020
Accepted: 08 Dec. 2020
Published: 04 Feb. 2021

How to cite this article:

Anderson CD, Makgotloe A, Koetsie K, Rose A. Survey of back and neck pain in South African ophthalmologists. *Afr Vision Eye Health*. 2021; 80(1), a616. <https://doi.org/10.4102/aveh.v80i1.616>

Copyright:

© 2021. The Author(s). Licensee: AOSIS. This work is licensed under the Creative Commons Attribution License.

Read online:



Scan this QR code with your smart phone or mobile device to read online.

Background: Back and neck pain are common in the general population and a common occupational condition. Occupationally related back and neck pain is common amongst ophthalmologists.

Aim: The aim of this study was to determine the prevalence of back and neck pain amongst South African ophthalmologists.

Setting: This was a survey that was conducted amongst South African ophthalmologists ($n = 324$) and ophthalmologists-in-training ($n = 115$).

Methods: The response rate was 44.8% ($n = 197$). There was no sampling and a hyperlink to an electronic questionnaire, hosted on the REDCap survey platform, was sent to 440 potential respondents. A descriptive analysis was carried out using STATA version 15. Ethical approval was obtained from the University of Witwatersrand.

Results: Mean age was 47.1 years (standard deviation [s.d.] ± 12.1) and 62 (31.5%) women. The cohort was stratified into specialists ($n = 81$, 41.1%), subspecialists ($n = 75$, 38.1%) and ophthalmologists-in-training ($n = 41$, 20.8%). Most of the participants ($n = 156$, 79.2%) were qualified ophthalmologists. Respondents reported current back pain ($n = 124$, 62.9%), lower back pain ($n = 66$, 33.5%) and neck pain ($n = 76$, 38.6%). The overall median pain rating (on a scale from 1 to 10) was 4. Respondents reported mild pain ($n = 71$, 36%) and severe pain ($n = 35$, 17.7%). Operating and using a slit lamp were pain aggravating factors in 36% ($n = 71$) and 37.1% ($n = 73$), respectively.

Conclusion: Back and neck pain are prevalent amongst South African ophthalmologists and comparable to other ophthalmologists worldwide.

Keywords: age; gender; height; job capacity; subspecialty; hours spent working; and specific questions about back and neck pain.

Introduction

Back and neck pain is widespread in our society today.^{1,2} The global prevalence of low back pain is estimated to be 38.9%.² The global prevalence of neck pain is much more varied because of the large variability within the studies which assess this factor. It is estimated to range between 0.4% and 86.8%.¹ This high prevalence rate is also true amongst surgeons.³ In the United States, 86.9% of laparoscopic surgeons report physical discomfort as a result of their work.⁴ In their systematic review that investigated the prevalence of work-related musculoskeletal disorders amongst surgeons and interventionalists ($n = 2815$), Epstein et al. (2017) found that the 12-months prevalence estimate of back and neck pain was 49% and 60%, respectively.³ Globally the prevalence of back and neck pain in ophthalmologists ranges between 26% – 82% and 18% – 93%, respectively.^{5,6,7,8,9,10,11,12,13,14}

Ophthalmology is a physically and mentally demanding speciality. Kitzmann et al. compared family physicians and eye care physicians and found that more eye care physicians classified themselves into a high strain job category and had more job factors associated with musculoskeletal symptoms than family physicians.¹⁰ Approximately half of ophthalmic plastic surgeons in a US-based survey agreed that specific components of their job could shorten their career length, interfere with their quality of life or lead to spinal disorders.¹² High patient loads and high volumes of surgery have been shown to contribute to the development of these symptoms.^{8,9} In a survey study of Saudi Arabian eye care professionals, musculoskeletal symptoms were reduced during vacation periods, thus suggesting a relationship with occupation.¹⁴ Ergonomics in ophthalmology are less than ideal. Shaw et al. investigated the mechanical exposure of vitreoretinal

surgeons and found that these ophthalmologists were often exposed to postures that are more demanding than neutral positions.¹⁵ Despite the ergonomic risks of the profession, many ophthalmologists are not aware of this problem. Of the ophthalmologists surveyed in India by Kaup et al., 27.9% ($n = 105/377$) had informed themselves on ergonomics and only 2.4% had used an ergonomist in their practice.⁶ In this study, there was a significant association between 'lack of good clinic and operation room ergonomic practices' and musculoskeletal symptoms.⁶

Musculoskeletal disorders may affect service delivery, and Dhimitri et al. found that 15% ($n = 697$) of ophthalmologists in their survey reported some limitation in their work.⁸ In a survey of ophthalmic plastic surgeons ($n = 130$), it was found that 54 (42%) surgeons had adjusted their operating practice, 10 (8%) had undergone surgery, and 12 (9%) had stopped operating as a result of spinal issues.¹²

Work-related musculoskeletal symptoms are widespread in ophthalmology, and the practice setting in which most ophthalmologists find themselves contributes to these problems.^{6,15} It has been recommended that improved ergonomics and education about these problems are important to reduce their impact.^{6,15} To the best of our knowledge back and neck pain in ophthalmologists has not been investigated in South Africa and this study aimed to address this knowledge gap. The aim of this study was to determine the prevalence of back and neck pain in South African ophthalmologists and associated aggravating factors.

Methods

Study population and data collection

This was a survey using an electronic questionnaire amongst South African based board-certified ophthalmologists and ophthalmologists-in-training. The study was based in South Africa. According to Resnikoff et al. and the International Council of Ophthalmology there are 324 registered ophthalmologists in the country.¹⁶ There are approximately 115 ophthalmologists-in-training in South Africa (this is an estimate from the Ophthalmological Society of South Africa, 2018). The ophthalmologists-in-training consisted of medical officers (pre-residents) and registrars (residents). The registrar (residency) programme is a four-year training programme. In South Africa, these are considered junior doctors. There was no sampling and all eligible doctors were invited to participate in the study. The questionnaire collected information on age, gender, height, job capacity, subspecialty, hours spent working and specific questions about back and neck pain. The questionnaire was administered on the REDCap platform, which is hosted by the University of Witwatersrand. This is a secure, web-based software platform designed for studies comprising of several research tools.^{17,18} The data were imported directly from this platform. The questionnaire was emailed to the participants who completed it anonymously. Informed consent was provided and participation was voluntary. Data were collected from

January to May 2019. Of the approximately 439 people invited, the cohort consisted of 197 people who completed the questionnaire giving a response rate of 44.8%.

Data analysis

A descriptive analysis was done using STATA version 15 (StataCorp, College Station, United States [US]). Frequencies were calculated for categorical data, mean and standard deviations (s.d.) were calculated for parametric data, medians and interquartile ranges (IQRs) were computed for non-parametric data. Odds ratios (ORs) with 95% confidence intervals (CIs) were calculated for risk factors related to back pain. A multivariate logistic analysis using a stepwise backward approach was performed to test the simultaneous association between back pain and different variables.

Ethical consideration

Ethics approval for the study was granted by the Human Research Ethics Committee (Medical) of the University of the Witwatersrand. (Certificate number: M180737).

Results

The age was normally distributed, and the mean age was 47.1 years (s.d. ± 12.1). The minimum age was 27 and the maximum age was 78. The age was categorised into two groups, < 47 years and ≥ 47 years, based on the mean age. There were 62 (31.5%) women and 135 (68.5%) men in the cohort.

The cohort was stratified into specialists ($n = 81$, 41.1%), subspecialists ($n = 75$, 38.1%) and junior doctors ($n = 41$, 20.8%). The majority of the participants ($n = 156$, 79.2%) were qualified ophthalmologists.

Figure 1 illustrates the proportion of self-reported areas of expertise of the 75 subspecialists. There were 19 (25.5%) specialists who indicated they worked in a subspecialty but had not indicated the specific speciality. In this category, we




FIGURE 1: The proportion of 75 subspecialists working in the various subcategories of ophthalmological specialties.

included participants who indicated they worked in more than one speciality.

The participants worked in the state sector ($n = 68$, 34.5%), private sector ($n = 108$, 54.8%) and a combination of state and the private sectors ($n = 21$, 10.7%). Table 1 describes the characteristics of the different strata and their occupational history. Participants self-reported the number of hours they worked per week in a clinic setting or performing surgery.

There were variable responses regarding reporting pain, how they subjectively quantified it and how they managed the pain. Participants were asked to report the pain in the previous month, the previous three months or in the previous year. They reported the factors that aggravated the pain. The pain was managed using either physiotherapy, medication, surgery and exercise or a combination of these modalities. From the cohort, the respondents who reported to be currently suffering from musculoskeletal back pain was 62.9% ($n = 124$). Lower back pain was reported by 33.5% of respondents ($n = 66$) and neck pain was reported by 38.6% ($n = 76$). The overall median pain rating (on a scale from 1 to 10) was 4. Most pain that was reported was

mild, 43.2% ($n = 85$), but severe pain was reported by 17.7% ($n = 35$). Operating and slit lamp utilisation were found to be aggravating factors for pain in 36% ($n = 71$) and 37.1% ($n = 73$), respectively. Table 2 describes the back and neck pain experienced by the doctors working in ophthalmology, pain characterisation, aggravating and relieving factors and treatment modalities.

Odds ratios were calculated for the association between various risk factors currently reporting muscular skeletal pain. These included age, sex, height, rank, trauma history, time spent working overall, time spent in the clinic weekly and time spent operating. None of these associations were found to be statistically significant. Table 3 describes these ORs.

A logistic regression model using a backward regression approach was used to assess covariates that predict for musculoskeletal pain. The following variables were assessed: age, sex, height, rank, time spent working overall, time spent in the clinic weekly and time spent operating. None of these associations were found to be statistically significant. Table 4 describes the covariates that predict for muscular skeletal pain.

TABLE 1: Demographics and occupational history of 197 doctors working in ophthalmology.

Covariate	Specialist $n = 81$ (41.1%)		Subspecialist $n = 75$ (38.1%)		Junior $n = 41$ (20.8%)		Total $N = 197$ (100.0%)	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Age in years								
Mean	49.9	-	51.6	-	33.0	-	47.1	-
s.d.	11.0	-	10.3	-	3.6	-	12.1	-
Minimum	31	-	34	-	27	-	27	-
Maximum	71	-	78	-	45	-	78	-
Sex								
Male	63	77.8	56	74.7	16	39.1	135	68.5
Female	18	22.2	19	25.3	25	60.9	62	31.5
Height (cm)								
Mean	173	-	177	-	170	-	174	-
s.d.	7.9	-	8.5	-	11.3	-	9.4	-
Work setting								
State	13	16.1	16	21.3	39	95.2	68	34.5
Private	62	76.5	45	60.0	1	2.4	108	54.8
Both	6	7.4	14	18.7	1	2.4	21	10.7
Hours worked								
Overall hours worked per week								
Median	40	-	40	-	39	-	38	-
IQR	32–42	-	34–43	-	30–42	-	26.5–50	-
≥ 40 h/week	71	87.7	59	78.7	30	73.2	160	81.2
< 40 h/week	10	12.3	16	21.3	11	26.8	37	18.8
Hours worked in clinic per week								
Median	30	-	25	-	30	-	30	-
IQR	24–35	-	20–30	-	20–32	-	22–33	-
≥ 30 h/week	53	65.4	29	38.7	22	53.7	104	52.8
< 30 h/week	28	34.6	46	61.3	19	46.3	93	47.2
Hours operating per week								
Median	8	-	10	-	6	-	8	-
IQR	6–10	-	8–15	-	3–8	-	3–8	-
≥ 8 h/week	54	66.7	57	76.0	17	41.5	120	64.9

Note: Doctors were stratified into three categories: specialists (81), subspecialists (75) and junior doctors (41).

IQR, interquartile range; s.d., standard deviation.

TABLE 2: Muscular skeletal pain experienced by the 197 doctors working in ophthalmology, the characterisation of their pain and the factors that aggravated and relieved the pain and the modalities they used to manage their pain.

Covariate	Specialist <i>n</i> = 81 (41.0%)		Subspecialist <i>n</i> = 75 (38.0%)		Junior <i>n</i> = 41 (21.0%)		Total <i>N</i> = 197 (100.0%)	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Current MS pain	54	66.7	46	61.3	24	58.5	124	62.9
Location of pain†								
Lower back	33	40.7	20	26.7	13	31.7	66	33.5
Mid back	8	9.9	4	5.3	6	14.6	18	9.1
Upper back	14	17.2	10	13.3	14	34.2	38	19.3
Neck	28	34.5	37	49.3	11	26.8	76	38.6
Pain rating‡								
Median	4	-	4	-	5	-	4	-
IQR	3–5.5	-	3–6	-	3–6	-	1–8	-
Category of pain								
No pain reported	29	35.8	29	38.7	19	46.3	77	39.1
Mild/moderate pain	39	48.1	33	44.0	13	31.7	85	43.2
Severe	13	16.1	13	17.3	9	22.0	35	17.7
History of trauma	9	11.1	18	24.0	1	2.4	28	14.2
Duration of pain in years								
Median	6	-	6	-	5	-	6	-
IQR	0–45	-	0–34	-	0–52	-	0–40	-
Subjective experience of pain reported by participants§								
In previous month	51	-	43	-	20	-	114	-
Median number of episodes reported	5	-	4	-	5	-	4	-
IQR	2–10	-	2–10	-	2.5–11	-	2–10	-
In previous 3 months	49	-	42	-	22	-	113	-
Median number of episodes reported	15	-	9.5	-	13.5	-	12	-
IQR	6–20	-	6–30	-	6–30	-	6–30	-
In previous year	47	-	40	-	23	-	110	-
Median number of episodes reported	40	-	30	-	30	-	30	-
IQR	12–100	-	12–76	-	12–240	-	12–96	-
Sequelae of pain								
Affects ADL	29	35.8	36	48.0	24	58.5	89	45.2
Resulted sick leave in past year	7	8.6	4	5.3	4	9.8	15	7.6
Median sick days	7	1–10	7	3–35	3.5	2–9	5	2–10
IQR	-	-	-	-	-	-	-	-
Aggravating factors								
Operating	26	32.1	34	45.3	11	26.8	71	36.0
Clinic work	14	17.3	5	6.7	7	17.1	26	13.2
Slit lamp utilisation	37	45.7	23	30.7	13	31.7	73	37.1
Pain management used								
Physiotherapy and cardiovascular	31	38.3	37	49.3	14	34.2	82	41.6
Surgery	1	1.2	3	4.0	0	-	4	2.0
Medication¶	37	45.7	22	29.3	13	31.7	72	36.6
Paracetamol	21	25.9	10	13.3	10	24.4	41	20.8
NSAID	33	40.7	20	26.7	11	-	64	32.5
Opiates	3	3.7	1	1.3	2	4.9	6	3.1
Exercise††	76	93.8	74	93.8	35	85.4	185	93.9
Cardio‡‡	57	70.4	59	78.7	29	70.7	145	73.6
Weights‡‡	19	23.5	20	26.7	11	26.8	50	25.4
Stretching‡‡	31	38.3	33	44.0	14	34.2	78	39.6

IQR, interquartile range; MS, musculoskeletal; ADL, activities of daily living; NSAID, nonsteroidal anti-inflammatory drug.

†, Participants may have indicated multiple areas that were affected.

‡, Pain was scored on a scale of 1–10.

§, The calculation is based on the number of participants indicating that they had experienced pain in the specified time periods.

¶, Participants may have indicated using more than one type of analgesia.

††, Exercised for at least 1 hour per week.

‡‡, May have participated in more than one type of exercise.

TABLE 3: Odds ratios and 95% confidence intervals for the association between various risk factors currently reporting muscular skeletal pain.

Covariate	OR	s.e.	95% CI	p
Age (years)				
≤ 47	0.726	0.215	0.406–1.298	0.280
> 47	-	-	-	-
Sex				
Females	1.876	0.627	0.975–3.611	0.060
Males	-	-	-	-
Height (centimetres)				
≤ 170	1.065	0.315	0.596–1.903	0.831
> 170	-	-	-	-
Category				
Specialists	1.265	0.453	0.627–2.553	0.512
Junior	-	-	-	-
Trauma				
No previous trauma	0.753	0.312	0.334–1.696	0.493
Previous trauma	-	-	-	-
Time spent working overall (h/week)				
≤ 40	0.834	0.311	0.401–1.732	0.627
> 40	-	-	-	-
Time spent in clinic (h/week)				
≤ 30	0.954	0.281	0.535–1.702	0.874
> 30	-	-	-	-
Time spent operating (h/week)				
≤ 8	1.731	0.555	0.924–3.244	0.087
> 8	-	-	-	-

Note: Standard error and *p*-values are reported as well.

OR, odds ratios; s.e., standard error; CI, confidence intervals.

TABLE 4: Logistic regression model using a backward regression approach.

Covariate	OR	s.e.	95% CI	p
Age	0.981	0.178	0.948–1.017	0.311
Sex	2.058	1.054	0.754–5.617	0.311
Height	1.012	0.022	0.969–1.057	0.585
Specialists	-	-	-	-
Subspecialist	0.756	0.287	0.359–1.592	0.287
Junior doctor	0.553	0.297	0.193–1.587	0.462
Hours per week	0.521	0.185	0.260–1.045	0.066
Clinic hours per week	1.015	0.185	0.260–1.045	0.066
Operating hour per week	1.013	0.332	0.950–1.081	0.684

Note: Covariates that predict for muscular skeletal disorder in ophthalmologists.

OR, odds ratios; s.e., standard error; CI, confidence intervals.

Discussion

This survey reported that 62.9% of the respondents suffered from back or neck pain. Compared with the global estimates of neck pain (mean 23.1%) the cohort had a higher prevalence of 38.6%.¹ The median pain score was 4, suggesting that although common, the pain perceived by these doctors is not severe. However, despite this, 45.2% of the entire cohort report that these symptoms had an effect on their activities of daily living. The cohort was predominantly male (68.5%), although, amongst the junior doctors the predominance was female (60.9%). This may suggest a change in the gender proportions of the ophthalmic community in the younger cohort. Male predominance in ophthalmology is common worldwide.^{7,8,9,11,13,19}

The prevalence of low back pain in this cohort (33.5%) compares most similarly to Canadian ophthalmologists (36%).¹¹ Neck pain prevalence in this cohort compares most similarly to ophthalmologists in India in a study by Venkatesh et al.: 38.6% versus 33%.⁹ The majority of these similar studies fell within a range of 26% – 50.6%^{5,6,7,8,9,10,11,12} for reported back pain and 18% – 58% for reported neck pain.^{5,6,7,8,9,10,11,12} Studies done in Iran and Saudi Arabia fell outside of this range; reporting back and neck pain to range between 80% – 82% and 69% – 93%, respectively.^{13,14} This comparison, therefore, suggests that South African ophthalmologists have comparable levels of reported neck and back pain to the rest of the world.

The median number of sick days taken was five days. Considering that the total number of working days per year is 260, these numbers are substantial. These self-reported symptoms have consequences beyond just the individual reporting them. From this cohort, four out of 197 respondents had surgery for musculoskeletal-related pain. Similarly, a United Kingdom-based study found that nine of 325 participating ophthalmologists required surgery for back pain.⁵ In a large systematic review of surgeons and interventionalists, 12% (*n* = 277/2319) of the physicians with work-related musculoskeletal disorders required leave, modified their practice or retired early.³ A survey of Australian optometrists found 35 reports of hospitalisation and 91 reports of changing jobs or duties because of work-related musculoskeletal symptoms.²⁰ It is thus clear that these self-reported symptoms can have an impact on service delivery and has consequences for ophthalmologists and their long-term quality of life.

The ergonomics of ophthalmic practice are challenging. In this study, 36% of survey respondents reported operating as an aggravating factor for musculoskeletal symptoms and 37.1% and 13.2% said that slit-lamp utilisation and clinic work, respectively, aggravated their symptoms. The median number of hours worked in the clinic per week was 30, and the median number of hours spent operating per week was 8; this suggests that South African ophthalmologists are frequently exposed to these aggravating factors in their workplace. Shaw et al. showed that vitreoretinal surgeons using the indirect ophthalmoscope were exposed to postures that were ergonomically undesirable.¹⁵ Work-related factors performed by eye care physicians such as repetitive tasks and awkward positions have been identified and shown to be a risk factor for musculoskeletal symptoms.^{10,20} The association between poor ergonomic practices and musculoskeletal symptoms has been demonstrated in ophthalmologists.⁶ We recommend that ophthalmologists would benefit from an ergonomic workplace assessment to mitigate developing musculoskeletal symptoms.

A strength of this research was as far as we could establish, that it was the first study in South Africa to look at neck and back pain in South African ophthalmologists and provided valuable insight into the topic in this group. Also, the response rate was good. The limitations of the study include that this was not a

random sample and thus may not be representative of the study population. Also, despite the good response rate the study may have been underpowered and had there been more respondents there may have been more statistically significant associations as were demonstrated in other similar studies. There may have been selection bias as ophthalmologists with back or neck pain may have opted into the study and recall bias may have been introduced in self-reported questions. The survey was limited to questions on musculoskeletal disorders related to the spine and other causes for these symptoms may have been missed. Workplace ergonomic assessments were not carried out and we recommend that future studies objectively assess musculoskeletal disorders in the workplace of the ophthalmologist and relate findings to self-reported symptoms. This may assist to inform recommendations for workplace adjustments.

Conclusion

In conclusion, back and neck pain are common amongst South African ophthalmologists, and they compare similarly to their global counterparts. The impact that these symptoms may have on ophthalmologists can be significant as it may affect their daily life and clinical practice. Ophthalmic practice is ergonomically challenging, and it is recommended that improving this in the workplace may assist in lowering the burden of these musculoskeletal complaints.

Acknowledgements

The authors would like to acknowledge all the doctors who participated in our study. We appreciate their time and effort.

Competing interests

The authors have declared that no competing interests exist.

Authors' contributions

C.D.A. assisted in the study conception. C.D.A., A.M. and K.K. were involved in study design and data collection. A.R. participated in data analysis. C.D.A., A.M., K.K. and A.R. drafted and reviewed the article. All authors reviewed and approved the final manuscript.

Funding information

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Data availability statement

Data are available on reasonable request from the corresponding author.

Disclaimer

The views and opinions expressed in this article are those of the authors and do not necessarily reflect the official policy or position of any affiliated agency of the authors.

References

- Hoy DG, Protani M, De R, Buchbinder R. The epidemiology of neck pain. *Best Pract Res Clin Rheumatol.* 2010;24(6):783–792. <https://doi.org/10.1016/j.berh.2011.01.019>
- Hoy D, Bain C, Williams G, et al. A systematic review of the global prevalence of low back pain. *Arthritis Rheum.* 2012;64(6):2028–2037. <https://doi.org/10.1002/art.34347>
- Epstein S, Sparer EH, Tran BN, et al. Prevalence of work-related musculoskeletal disorders among surgeons and interventionalists. *JAMA Surg.* 2018;153(2):e174947. <https://doi.org/10.1001/jamasurg.2017.4947>
- Park A, Lee G, Seagull FJ, Meenaghan N, Dexter D. Patients benefit while surgeons suffer: An impending epidemic. *J Am Coll Surg.* 2010;210(3):306–313. <https://doi.org/10.1016/j.jamcollsurg.2009.10.017>
- Chatterjee A, Ryan WG, Rosen ES. Back pain in ophthalmologists. *Eye.* 1994; 8(4):473–474. <https://doi.org/10.1038/eye.1994.112>
- Kaup S, Shivalli S, Kulkarni U, Arunachalam C. Ergonomic practices and musculoskeletal disorders among ophthalmologists in India: An online appraisal. *Eur J Ophthalmol.* 2020;30(1):196–200. <https://doi.org/10.1177/1120672118815107>
- Hyer JN, Lee RM, Chowdhury HR, Smith HB, Dhital A, Khandwala M. National survey of back & neck pain amongst consultant ophthalmologists in the United Kingdom. *Int Ophthalmol.* 2015;35(6):769–775. <https://doi.org/10.1007/s10792-015-0036-z>
- Dhimitri KC, McGwin G, McNeal SF, et al. Symptoms of musculoskeletal disorders in ophthalmologists. *Am J Ophthalmol.* 2005;139(1):179–181. <https://doi.org/10.1016/j.ajo.2004.06.091>
- Venkatesh R, Kumar S. Back pain in ophthalmology: National survey of Indian ophthalmologists. *Indian J Ophthalmol.* 2017;65(8):678–682. https://doi.org/10.4103/ijo.IJO_344_17
- Kitzmann AS, Fethke NB, Baratz KH, Zimmerman MB, Hackbarth DJ, Gehrs KM. A survey study of musculoskeletal disorders among eye care physicians compared with family medicine physicians. *Ophthalmol.* 2012;119(2):213–220. <https://doi.org/10.1016/j.ophtha.2011.06.034>
- Diaconita V, Uhlman K, Mao A, Mather R. Survey of occupational musculoskeletal pain and injury in Canadian ophthalmology. *Can J Ophthalmol.* 2019;54(3):314–322. <https://doi.org/10.1016/j.cjco.2018.06.021>
- Sivak-Callcott JA, Diaz SR, Ducatman AM, Rosen CL, Nimbar AD, Sedgeman JA. A survey study of occupational pain and injury in ophthalmic plastic surgeons. *Ophthalmic Plast Reconstr Surg.* 2011;27(1):28–32. <https://doi.org/10.1097/IOP.0b013e3181e99cc8>
- Chams H, Mohammadi SF, Moayeri A. Frequency and assortment of self-reported occupational complaints among Iranian ophthalmologists: A preliminary survey. *Med Gen Med.* 2004;6(4):1.
- Al-Marwani Al-Juhani M, Khandekar R, Al-Harby M, Al-Hassan A, Edward DP. Neck and upper back pain among eye care professionals. *Occup Med.* 2015; 65(9):754–757. <https://doi.org/10.1093/occmed/kqv132>
- Shaw C, Bourkiza R, Wickham L, McCarthy I, Mckechnie C. Mechanical exposure of ophthalmic surgeons: A quantitative ergonomic evaluation of indirect ophthalmoscopy and slit-lamp biomicroscopy. *Can J Ophthalmol.* 2017; 52(3):302–307. <https://doi.org/10.1016/j.cjco.2016.09.011>
- Resnikoff S, Felch W, Gauthier TM, Spivey B. The number of ophthalmologists in practice and training worldwide: A growing gap despite more than 200 000 practitioners. *Br J Ophthalmol.* 2012;96(6):783–787. <https://doi.org/10.1136/bjophthalmol-2011-301378>
- Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap) – A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform.* 2009;42(2):377–381. <https://doi.org/10.1016/j.jbi.2008.08.010>
- Harris PA, Taylor R, Minor BL, et al. The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform.* 2019;95(December 2018):103208. <https://doi.org/10.1016/j.jbi.2019.103208>
- McAlister C, Jin YP, Braga-Mele R, DesMarchais BF, Buys YM. Comparison of lifestyle and practice patterns between male and female Canadian ophthalmologists. *Can J Ophthalmol.* 2014;49(3):287–290. <https://doi.org/10.1016/j.cjco.2014.02.007>
- Long J, Naduvilath TJ, Hao L, et al. Risk factors for physical discomfort in Australian optometrists. *Optom Vis Sci.* 2011;88(2):317–326. <https://doi.org/10.1097/OPX.0b013e3182045a8e>